

Recombinant Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody

Rabbit Monoclonal Antibody [Clone CDH16/1532R]

Catalog No	Format	Size
1014-RBM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1014-RBM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1014-RBM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	CDH16/1532R
Gene Name	CDH16
Immunogen	Recombinant human full-length CDH16 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	130kDa
Cellular Localization	Cell membrane
Species Reactivity	Guinea Pig, Hamster, Human, Mouse, Rabbit, Rat
Positive Control	Normal kidney or renal cell carcinoma. Kidney.

*Optimal dilution for a specific application should be determined.



Product Images for Recombinant Ksp-Cadherin (Kidney-Specific Cadherin) / CDH16 Antibody



Formalin-fixed, paraffin-embedded Rat Kidney stained with KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)





Formalin-fixed, paraffin-embedded Mouse Kidney stained with KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)




SDS-PAGE Analysis of Purified KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)

Flow Cytometric Analysis of Human HEK293 cells using KSP-Cadherin Recombinant Rabbit Monoclonal Antibody (CDH16/1532R) followed by Goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).



Western Blot analysis of Human Kidney tissue lysate using CDH16 Recombinant Rabbit Monoclonal Antibody (CDH16/1532R).

Western Blot Analysis of Human Kidney, Mouse Kidney, Rat Kidney, Hamster Kidney, Guinea pig Kidney and Rabbit Kidney tissue lysates using Ksp-Cadherin Recombinant Rabbit Monoclonal Antibody (CDH16/1532R).



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with KSP-Cadherin Rabbit Recombinant Monoclonal Antibody (CDH16/1532R)

Specificity & Comments

This MAb recognizes a protein of 130kDa, identified as Ksp-cadherin. Cadherins form a superfamily of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins have been implicated in embryogenesis, tissue morphogenesis, tissue structure maintenance, cell polarization, neoplastic invasiveness and metastasis, and membrane transport. It is suggested that Ksp-cadherin is a marker for terminal differentiation of the basolateral membranes of renal tubular epithelial cells. Within the kidney, Ksp-Cadherin is found exclusively in the basolateral membrane of renal tubular epithelial cells and collecting duct cells, and not in glomeruli, renal interstitial cells, or blood vessels. Ksp-Cadherin has been suggested to distinguish Chromophobe Renal-Cell Carcinoma from Oncocytoma.

Supplied As

200ug/ml of Ab Purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.