

p21WAF1 (Tumor Suppressor Protein) Antibody

Mouse Monoclonal Antibody [Clone CIP1/823]


Catalog No	Format	Size
1026-MSM3-P0	Purified Ab with BSA and Azide	200ug/ml
1026-MSM3-P1	Purified Ab with BSA and Azide	200ug/ml
1026-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunofluorescence (IF)	1-3ug/ml
Immunohistochemistry (IHC)	1-2ug/ml
Western Blot (WB)	2-4ug/ml


Product Details	
Clone	CIP1/823
Gene Name	CDKN1A
Immunogen	Recombinant full-length human CDKN1A protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	21kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human
Positive Control	colon or breast carcinoma (IHC)., HeLa cells (IF). HeLa cell lysates (WB). Skin

**Optimal dilution for a specific application should be determined.*


Product Images for p21WAF1 (Tumor Suppressor Protein) Antibody




Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with p21 Mouse Monoclonal Antibody (CIP1/823).



Western Blot Analysis of human HeLa cell lysate using p21 Mouse Monoclonal Antibody (CIP1/823).



Immunofluorescence Analysis of HeLa cells labeling p21 with p21 Mouse Monoclonal Antibody (CIP1/823) followed by Goat anti-Mouse IgG-CF555 (Red). Membrane stained with Phalloidin 488 (Green)



Formalin-fixed, paraffin-embedded human Lung SqCC stained with p21 Mouse Monoclonal Antibody (CIP1/823).

Specificity & Comments

This MAb recognizes a 21kDa protein, identified as the p21WAF1 tumor suppressor protein. This MAb is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdk's and a tumor suppressor involved in the pathogenesis of a variety of malignancies. The expression of this gene acts as an inhibitor of the cell cycle during G1 phase and is tightly controlled by the tumor suppressor protein p53. Its expression is induced by the wild type, but not mutant, p53 suppressor protein. Normal cells generally display a rather intense nuclear p21 expression. Loss of p21 expression has been reported in many carcinomas (gastric carcinoma, non-small cell lung carcinoma, thyroid carcinoma).

Research Areas

AKT Signaling, Bladder Cancer, Breast Cancer, Cancer, Cardiovascular, Colon Cancer, Cytokine Signaling, Developmental Biology, Immunology, Infectious Disease, Lung Cancer, Nuclear Marker, Ovarian Cancer, Signal Transduction, Transcription Factors

Known Applications & Suggested Dilutions

Immunofluorescence (1-2ug/ml) | Western Blot (1-2ug/ml) | Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes) | Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.