

p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody

Mouse Monoclonal Antibody [Clone KIP1/1357]


Catalog No	Format	Size
1027-MSM7-P0	Purified Ab with BSA and Azide	200ug/ml
1027-MSM7-P1	Purified Ab with BSA and Azide	200ug/ml
1027-MSM7-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml


Product Details	
Clone	KIP1/1357
Gene Name	CDKN1B
Immunogen	Recombinant full-length human CDKN1B protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	25-26kDa
Cellular Localization	Cytoplasm, Endosome, Nucleus
Species Reactivity	Human
Positive Control	Breast, Cervical or Colon Carcinoma., HeLa or MCF7 cells. Human tonsil

**Optimal dilution for a specific application should be determined.*


Product Images for p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody




IHC analysis of formalin-fixed, paraffin-embedded human prostate. Strong nuclear staining using KIP1/1357 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



IHC analysis of formalin-fixed, paraffin-embedded human brain. Strong nuclear staining using KIP1/1357 at 2ug/ml in PBS for 30min RT. Inset: PBS instead of primary antibody; secondary only negative control.



IHC analysis of formalin-fixed, paraffin-embedded human colon. Strong nuclear staining using KIP1/1357 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified p27 Mouse Monoclonal Antibody (KIP1/1357). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

This MAbs recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human breast cancer cell lines (ZR75-1, ZR75-30, MCF-7, MDAMB453, T47D, CAL51, 734B), the antibody labels a single band corresponding to p27Kip1. It functions as a negative regulator of G1 progression and has been proposed to function as a possible mediator of TGF- β induced G1 arrest. p27Kip1 is a candidate tumor suppressor gene. Reportedly, low p27 expression has been associated with unfavorable prognosis in renal cell carcinoma, colon carcinoma, breast carcinomas, non-small-cell lung carcinoma, hepatocellular carcinoma, multiple myeloma, and lymph node metastases in papillary carcinoma of the thyroid, as well as a more aggressive phenotype in carcinoma of the cervix.

Research Areas

Cardiovascular, Immunology, AKT Signaling, Cytokine Signaling, Infectious Disease, Lung Cancer, Nuclear Marker, Ovarian Cancer, Signal Transduction, Transcription Factors

Known Applications & Suggested Dilutions

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.