

Carcinoembryonic Antigen (CEA) / CD66 Antibody

Mouse Monoclonal Antibody [Clone C66/1292]

| Catalog No | Format | Size |
|------------------|-----------------------------------|----------|
| 1048-MSM12-P0 | Purified Ab with BSA and Azide | 200ug/ml |
| 1048-MSM12-P1 | Purified Ab with BSA and Azide | 200ug/ml |
| 1048-MSM12-P1ABX | Purified Ab WITHOUT BSA and Azide | 1.0mg/ml |

| Applications | Tested Dillution |
|-------------------------|---------------------|
| Flow Cytometry (Flow) | 1-2ug/million cells |
| Immunofluorescence (IF) | 1-3ug/ml |

| Product Details | |
|------------------------|---|
| Clone | C66/1292 |
| Gene Name | CEACAM5 |
| Immunogen | Recombinant full-length human CEA protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2b |
| Mol. Weight of Antigen | 80-200kDa |
| Cellular Localization | Apical cell membrane, Cell membrane, Cell surface |
| Species Reactivity | Human |
| Positive Control | MCF7 or 293T cells. Human colon carcinoma. |

**Optimal dilution for a specific application should be determined.*

Product Images for Carcinoembryonic Antigen (CEA) / CD66 Antibody

Specificity & Comments

This antibody recognizes proteins of 80-200kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MAb does not react with nonspecific cross-reacting antigen (NCA) and with human polymorphonuclear leucocytes. It shows no reaction with a variety of normal tissues and is suitable for staining of formalin/paraffin tissues. CEA is not found in benign glands, stroma, or malignant prostatic cells. Antibody to CEA is useful in detecting early foci of gastric carcinoma and in distinguishing pulmonary adenocarcinomas (60-70% are CEA+) from pleural mesotheliomas (rarely or weakly CEA+). Anti-CEA positivity is seen in adenocarcinomas from the lung, colon, stomach, esophagus, pancreas, gallbadder, urachus, salivary gland, ovary, and endocervix.

Research Areas

Hematopoietic Stem Cells

Known Applications & Suggested Dilutions

Flow Cytometry (0.5-1ug/million cells) | Immunofluorescence (1-2ug/ml), Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.