

DMRT2 (Transcription Factor) Antibody

Mouse Monoclonal Antibody [Clone PCR-P-DMRT2-1B11]


Catalog No	Format	Size
10655-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10655-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10655-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Western Blot (WB)	2-4ug/ml	


Product Details	
Clone	PCR-P-DMRT2-1B11
Gene Name	DMRT2
Immunogen	Recombinant full-length human DMRT2 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b
Mol. Weight of Antigen	62kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	HeLa or MCF-7 cells.

*Optimal dilution for a specific application should be determined.


Product Images for DMRT2 (Transcription Factor) Antibody



SDS-PAGE Analysis. Purified DMRT2 Mouse Monoclonal Antibody (PCR-P-DMRT2-1B11). Confirmation of Integrity and Purity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using DMRT2 Mouse Monoclonal Antibody (PCR-P-DMRT2-1D3). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Flow Cytometric Analysis of PFA-fixed HeLa cells. DMRT2 Mouse Monoclonal Antibody (PCRP-DMRT2-1B11) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).

Specificity & Comments

The protein encoded by this gene belongs to the DMRT gene family, sharing a DM DNA-binding domain with *Drosophila* 'doublesex' (dsx) and *C. elegans* mab3, genes involved in sex determination in these organisms. Also, this gene is located in a region of the human genome (chromosome 9p24.3) associated with gonadal dysgenesis and XY sex reversal. Hence this gene is one of the candidates for sex-determining gene(s) on chr 9. Transcriptional activator that directly regulates early activation of the myogenic determination gene MYF5 by binding in a sequence-specific manner to the early epaxial enhancer element of it. Involved in somitogenesis during embryogenesis and somite development and differentiation into sclerotome and dermomyotome. Required for the initiation and/or maintenance of proper organization of the sclerotome, dermomyotome and myotome (By similarity).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.