Recombinant CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Antibody

Rabbit Monoclonal Antibody [Clone CFTR/1775R]

Catalog No	Format	Size
1080-RBM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1080-RBM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1080-RBM5-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	CFTR/1775R
Immunogen	Recombinant fragment (around aa590-830) of the human CFTR protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	168.14kDa
Cellular Localization	Apical cell membrane, Cell membrane, Early endosome membrane, Endoplasmic reticulum membrane, Nucleus, Recycling endosome membrane
Species Reactivity	Human
Positive Control	Kidney or Placenta., MOLT-4 cells. Pancreas

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant CFTR (Cystic Fibrosis Transmembrane Conductance Regulator) Antibody



SDS-PAGE Analysis of Purified Cystic fibrosis transmembrane conductance regulator Recombinant Rabbit Monoclonal Antibody (CFTR/1775R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Pancreas stained with CFTR Rabbit Recombinant Monoclonal Antibody (CFTR/1775R).

Specificity & Comments

Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts. CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.