

LH, alpha (Luteinizing Hormone, alpha) Antibody

Mouse Monoclonal Antibody [Clone LHα/756]

Catalog No	Format	Size
1081-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1081-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1081-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	LHα/756
Gene Name	CGA
Immunogen	Recombinant full-length hLH alpha protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	~13kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Anterior Pituitary.

*Optimal dilution for a specific application should be determined.

Product Images for LH, alpha (Luteinizing Hormone, alpha) Antibody



SDS-PAGE Analysis of Purified LH-alpha Monoclonal Mouse Antibody(LHα/756). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human Pituitary stained with LH alpha Mouse Monoclonal Antibody (LHα/756).

Specificity & Comments

This MAb reacts with a protein of ~13kDa, identified as alpha subunit of Luteinizing Hormone (LH). Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimulating hormone (TSH), and human chorionic gonadotropin (hCG). The protein dimer contains 2 polypeptide units, labeled alpha and beta subunits that are connected by two bridges. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids. The beta subunits vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action and is responsible for interaction with the LH receptor. This beta subunit contains the same amino acids in sequence as the beta subunit of hCG and both stimulate the same receptor; however, the hCG beta subunit contains an additional 24 amino acids and the hormones differ in the composition of their sugar moieties. LH is synthesized and secreted by gonadotrophs in the anterior lobe of the pituitary gland. In concert with the other pituitary gonadotropin follicle-stimulating hormone (FSH), it is necessary for proper reproductive function. In the female, an acute rise of LH levels triggers ovulation. In the male, where LH has also been called Interstitial Cell-Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. LH is a useful marker in classification of pituitary tumors and the study of pituitary disease.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Signal Transduction, Transcription Factors