

CD268 / BAFFR / TNFRSF13C Antibody

Mouse Monoclonal Antibody [Clone BAFFR/1558]

Catalog No	Format	Size
115650-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
115650-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
115650-MSM2-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	BAFFR/1558
Gene Name	TNFRSF13C
Immunogen	Recombinant full-length human CD268 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	19kDa (Monomer); 40kDa (Dimer)
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	Raji cells. Tonsil and lymph node.

*Optimal dilution for a specific application should be determined.

Product Images for CD268 / BAFFR / TNFRSF13C Antibody



SDS-PAGE Analysis of Purified CD268 / BAFFR Mouse Monoclonal Antibody (BAFFR/1558). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human tonsil stained with CD268 / BAFFR Mouse Monoclonal Antibody (BAFFR/1558).

Specificity & Comments

Defects in TNFRSF13C are the cause of immunodeficiency common variable type 4 (CVID4) [MIM:613494]; also called antibody deficiency due to BAFFR defect. CVID4 is a primary immunodeficiency characterized by antibody deficiency, hypogammaglobulinemia, recurrent bacterial infections and an inability to mount an antibody response to antigen. The defect results from a failure of B-cell differentiation and impaired secretion of immunoglobulins; the numbers of circulating B cells is usually in the normal range, but can be low.

Supplied As

200ug/ml of Ab purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology, AKT Signaling, B Cell Markers, Cytokine Signaling

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.