

Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) Antibody

Mouse Monoclonal Antibody [Clone SERPINA3/4189]

Catalog No	Format	Size
12-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
12-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
12-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	SERPINA3/4189
Gene Name	SERPINA3
Immunogen	Recombinant human Antichymotrypsin (AACT) protein fragment (around aa 49-187) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	65-76kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	HeLa cells. Human tonsil, pancreas or histiocytoma.

*Optimal dilution for a specific application should be determined.


Product Images for Alpha-1-Antichymotrypsin (SERPINA3) (Histiocytoma Marker) Antibody



SDS-PAGE Analysis of Purified AACT Mouse Monoclonal Antibody (SERPINA3/4189). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing >19,000 full-length human proteins using Monospecific to Alpha-1-Antichymotrypsin Mouse Monoclonal Antibody (SERPINA3/4189). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human liver stained with Alpha-1-Antichymotrypsin Mouse Monoclonal Antibody (SERPINA3/4189).

Specificity & Comments

It recognizes a protein of 65-76kDa, which is identified antichymotrypsin (AAT). AAT is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AAT is about one-tenth that of α -1-antitrypsin (AAT), with which it shares nucleic acid and protein sequence homology. Both are major acute phase reactants; their concentrations in plasma increase in response to trauma, surgery and infection. Elevated levels of AAT are widely, but not universally, reported in the cerebrospinal fluid and plasma of AD patients. Prostate-specific antigen (PSA) and its SDS-stable complex with AAT are in widespread use as markers for the diagnosis of prostate cancer. AAT deficiency may also be a possible cause of chronic liver disease. AAT antibody reacts with histiocytes and histiocytic neoplasms. It is widely used to identify histiocytes and tumors derived from them. Acinar tumors of the pancreas and salivary gland may also exhibit AAT positivity.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology