

## CD35 / CR1 (Follicular Dendritic Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone To5]

| Catalog No      | Format  | Size   |
|-----------------|---|--------|
| 1378-MSM3-P0    | Purified Ab with BSA and Azide at 200ug/ml    | 20 ug  |
| 1378-MSM3-P1    | Purified Ab with BSA and Azide at 200ug/ml    | 100 ug |
| 1378-MSM3-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |



| Applications               | Tested Dillution    | Note  |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow)      | 1-2ug/million cells |   |
| Immunofluorescence (IF)    | 1-3ug/ml            |   |
| Immunohistochemistry (IHC) | 1-2ug/ml            | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

### Product Details

|                               |   |
|-------------------------------|---|
| <b>Clone</b>                  | To5   |
| <b>Gene Name</b>              | CR1   |
| <b>Immunogen</b>              | Recombinant human CR1 protein               |
| <b>Host</b>                   | Mouse                                       |
| <b>Clonality</b>              | Monoclonal                                  |
| <b>Isotype / Light Chain</b>  | IgG1 / Kappa                                |
| <b>Mol. Weight of Antigen</b> | 210-220kDa                                  |
| <b>Cellular Localization</b>  | Membrane                                    |
| <b>Species Reactivity</b>     | Human                                       |
| <b>Positive Control</b>       | Follicular dendritic cells (FDC) in tonsil. |

\*Optimal dilution for a specific application should be determined.

### Product Images for CD35 / CR1 (Follicular Dendritic Cell Marker) Antibody



SDS-PAGE Analysis Purified CD35 Mouse Monoclonal Antibody (To5).  
Confirmation of Integrity and Purity of Antibody.

Formalin-fixed, paraffin-embedded human Tonsil stained with CD35 Mouse Monoclonal Antibody (To5).

**Specificity & Comments**

Recognizes a protein of 210-220kDa, which is identified as the complement receptor 1 (CR1)/CD35. This MAb does not block CR1 activity. It is highly specific to CR1 and shows no cross-reaction with CR2. The primary function of CR1 is to serve as the cellular receptor for C3b and C4b, the most important components of the complement system leading to clearance of foreign macromolecules. The Knops blood group system is a system of antigens located on this protein. Follicular dendritic cells (FDC) are restricted to the B-cell regions of secondary lymphoid follicles. They are CD21+/CD35+/CD1a-. This MAb labels follicular dendritic cells and follicular dendritic cell sarcoma.

---

**Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

**Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

**Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

**Research Areas**

Complement System, Immunology, Transcription Factors

---