

AIF1 / Iba1 (Microglia Marker) Antibody

Mouse Monoclonal Antibody [Clone AIF1/1909]

| Catalog No | Format | Size |
|----------------|-----------------------------------------------|--------|
| 199-MSM1-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 199-MSM1-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 199-MSM1-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details


| | |
|-------------------------------|-------------------------------------------------------------------------------------------------------|
| Clone | AIF1/1909 |
| Gene Name | AIF1 |
| Immunogen | Purified fragment of human recombinant AIF1 protein (around aa 1-146) (exact sequence is proprietary) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 17kDa |
| Cellular Localization | Cell projection, Cytoplasm, Cytoskeleton, Phagocytic cup, Ruffle membrane |
| Species Reactivity | Human |
| Positive Control | Human lymph node, tonsil or kidney. |

*Optimal dilution for a specific application should be determined.


Product Images for AIF1 / Iba1 (Microglia Marker) Antibody




SDS-PAGE Analysis of Purified AIF1 / Iba1 Mouse Monoclonal Antibody (AIF1/1909). Confirmation of Integrity and Purity of Antibody.




Formalin-fixed, paraffin-embedded human Lymph Node stained with AIF1 /Iba1 Mouse Monoclonal Antibody (AIF1/1909).



Formalin-fixed, paraffin-embedded human Kidney stained with AIF1 / Iba1 Mouse Monoclonal Antibody (AIF1/1909).



Analysis of Protein Array containing more than 19,000 full-length human proteins using AIF1 Mouse Monoclonal Antibody (AIF1/1909) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Tonsil stained with AIF1 / Iba1 Mouse Monoclonal Antibody (AIF1/1909).

Specificity & Comments

AIF1 is a cytoplasmic, calcium-binding protein that is thought to play a role in macrophage activation and function. AIF1, containing two EF domains, is induced by cytokines and Interferons. In an unstimulated state, AIF1 colocalizes with actin, and upon stimulation, translocates to lamellipodia. It is also a marker of human microglia and is expressed by macrophages in injured skeletal muscle. The gene encoding AIF1 resides in the tumor necrosis factor (TNF) cluster of genes, located in the region represented by the human major histocompatibility complex (MHC).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Cellular Markers and Tags, Dendritic Cell Marker