

## Factor XIIIa (Coagulation Factor XIIIa Chain) Antibody

Mouse Monoclonal Antibody [Clone F13A1/1448]

Catalog No	Format	Size
2162-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2162-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2162-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	F13A1/1448
<b>Gene Name</b>	F13A1
<b>Immunogen</b>	Recombinant fragment of human Factor XIIIa protein (aa46-181) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	83kDa
<b>Cellular Localization</b>	Cytoplasm, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	293T cells. Placenta.

\*Optimal dilution for a specific application should be determined.



### Product Images for Factor XIIIa (Coagulation Factor XIIIa Chain) Antibody



Western Blot Analysis of Recombinant Protein and HeLa cell lysate using Factor XIIIa Mouse Monoclonal Antibody (F13A1/1448).



SDS-PAGE Analysis of Purified Factor XIIIa Mouse Monoclonal Antibody (F13A1/1448). Confirmation of Integrity and Purity of Antibody.



Formalin-fixed, paraffin-embedded human Histiocytoma stained with Factor XIIIa Mouse Monoclonal Antibody (F13A1/1448).

Analysis of Protein Array containing more than 19,000 full-length human proteins using Factor XIIIa Mouse Monoclonal Antibody (F13A1/1448) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

### Specificity & Comments

The specificity of this monoclonal antibody to its intended target was validated by HuProt™ Array, containing more than 19,000, full-length human proteins. It recognizes a protein of 83kDa, which is identified as Factor XIIIa. It has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macrophages and dermal dendritic cells. Anti-factor XIIIa has been found to be useful in differentiating between dermatofibroma (almost all cases are positive), dermatofibrosarcoma protuberans (-/+) and desmoplastic malignant melanoma (-). Anti-factor XIIIa positivity is also seen in capillary hemangioblastoma, hemangioendothelioma, hemangiopericytoma, xanthogranuloma, xanthoma, hepatocellular carcinoma, glomus tumor, and meningioma.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Complement System, Cytokine Signaling, Immunology