

ICOS-L / ICOS Ligand / B7RP-1 (Immuno-Oncology Target) Antibody

Mouse Monoclonal Antibody [Clone ICOSL/3111]


Catalog No	Format	Size
23308-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
23308-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
23308-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes


Product Details	
Clone	ICOSL/3111
Gene Name	ICOSLG
Immunogen	Recombinant fragment of human ICOSLG (B7RP-1) protein (around aa 23-149) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	36kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	HL60, HuT78 and HUVEC lysates (WB). Human tonsil tissue (IHC)., U937

**Optimal dilution for a specific application should be determined.*


Product Images for ICOS-L / ICOS Ligand / B7RP-1 (Immuno-Oncology Target) Antibody




Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with ICOS-L Mouse Monoclonal Antibody (ICOSL/3111).




Flow Cytometric Analysis of U937 cells using ICOS-L Mouse Monoclonal Antibody (ICOSL/3111) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Immunofluorescence staining of U937 cells using ICOS-L Mouse Monoclonal Antibody (ICOSL/3111) followed by goat anti-mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot



SDS-PAGE Analysis Purified ICOS-L Mouse Monoclonal Antibody (ICOSL/3111). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using ICOS-L Mouse Monoclonal Antibody (ICOSL/3111). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with ICOS-L Mouse Monoclonal Antibody (ICOSL/3111).

Specificity & Comments

This MAb recognizes a protein of 36kDa, identified as ICOS-L. It is a ligand for the ICOS receptor that initiates T and B cell proliferation and cytokine secretion. ICOS-L interactions play an essential role in T cell-dependent B cell activation in peripheral lymphoid organs such as spleen and lymph nodes. ICOS-L protein is present in myeloid leukocytes, and by Northern blot there are 2.4, 3.0, and 7.0 kb transcripts in brain, heart, kidney, and liver, with lower expression in colon and thymus, and a 1.1 kb transcript in leukocytes. Tumor necrosis factor alpha (TNF), granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin-4 (IL-4) enhance its expression. LPS-induced up-regulation of ICOS-L is dependent on the MyD88-dependent signaling pathway.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology