

Fibronectin (8th Type III Repeat) Antibody

Mouse Monoclonal Antibody [Clone 568]

Catalog No	Format	Size
2335-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2335-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2335-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	568
Gene Name	Fn1
Immunogen	High molecular weight proteins secreted by cultivated human fibroblasts
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	220kDa (monomer); 440kDa (dimer)
Cellular Localization	Extracellular matrix, Extracellular space, Secreted
Species Reactivity	Human
Positive Control	SW156 cells. Human Kidney, Human Placenta

*Optimal dilution for a specific application should be determined.


Product Images for Fibronectin (8th Type III Repeat) Antibody




SDS-PAGE Analysis of Purified Fibronectin Monoclonal Antibody (568). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Placenta tissue lysate using Fibronectin Mouse Monoclonal Antibody (568).



Western Blot Analysis of Human Placenta tissue lysate using Fibronectin Mouse Monoclonal Antibody (568).



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Fibronectin Monoclonal Antibody (568).

Specificity & Comments

Epitope of this MAb is located in the 8th type III repeat in the cell-binding region of fibronectin. Fibronectins are disulfide-linked, dimeric glycoproteins of ~440kDa. They possess at least four binding sites for collagen, glycosaminoglycans, transglutaminase, and a cell surface receptor. Fibronectins are involved in cell adhesion, tissue organization, and wound healing. Fibronectins are present in basement membranes, interstitial connective tissue matrix, and blood. Cellular fibronectin is widely distributed in the stroma of many malignant tumors. This MAb is excellent for staining of formalin-fixed, paraffin-embedded tissues.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Articular Cartilage Extracellular Matrix, Cytokine Signaling, Infectious Disease, Lung Cancer, Mesenchymal Stem Cell Differentiation, Signal Transduction

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.