

GDF9 (Growth Differentiation Factor 9) Antibody

Mouse Monoclonal Antibody [Clone GDF9/4261]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 2661-MSM1-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 2661-MSM1-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 2661-MSM1-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |



| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details

| | |
|------------------------|---|
| Clone | GDF9/4261 |
| Gene Name | GDF9 |
| Immunogen | Tuberculin coupled peptide with sequence VPAKYSPLSVLTIEPDGSIAYKEYEDMIATKC that recognizes an epitope with the EPDG sequence near the C-terminal region of human GDF9. |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 |
| Mol. Weight of Antigen | 51kDa |
| Cellular Localization | Secreted |
| Species Reactivity | Human |
| Positive Control | Human ovary. |

*Optimal dilution for a specific application should be determined.

Product Images for GDF9 (Growth Differentiation Factor 9) Antibody



Formalin-fixed, paraffin-embedded human ovary stained with GDF9 Mouse Monoclonal Antibody (GDF9/4261).

SDS-PAGE Analysis of Purified GDF9 Mouse Monoclonal Antibody (GDF9/4261) Confirmation of Integrity and Purity of Antibody.

Specificity & Comments

GDF9 is a member of the bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. Growth factors synthesized by ovarian somatic cells directly affect oocyte growth and function. GDF9 is expressed in oocytes and is thought to be required for ovarian folliculogenesis. GDF9/4261 can be used in assays to detect oocyte expression and has been shown to neutralize GDF9 biological activity.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.