

Recombinant PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target) Antibody

Rabbit Monoclonal Antibody [Clone PDL1/4451R]


Catalog No	Format	Size
29126-RBM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
29126-RBM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
29126-RBM11-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


Product Details	
Clone	PDL1/4451R
Gene Name	CD274
Immunogen	Recombinant fragment (around aa 39-191) of human CD274 (PD-L1) protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	37-50kDa
Cellular Localization	Cell membrane, Early endosome membrane, Endomembrane system, Recycling endosome membrane
Species Reactivity	Human, Mouse
Positive Control	Human heart, Placenta, RAW or HepG2 cells., spleen or tonsil tissue. Jurkat, A549, MDA-MB-231, Raw.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant PD-L1 / PDCD1LG1 / CD274 / B7-H1 (Cancer Immunotherapy Target) Antibody




Formalin-fixed, paraffin-embedded human Lung SqCC stained with PD-L1 Recombinant Rabbit Monoclonal Antibody (PDL1/4451R).



Western Blot Analysis of A431, MCF-7, JEG-3, MDA-MB231, SW732 cell lysates using PD-L1 Recombinant Rabbit Monoclonal Antibody (PDL1/4451R).



Western Blot Analysis of A549, MDA-MB231 and Raw cell lysates using PD-L1 Recombinant Rabbit Monoclonal Antibody (PDL1/4451R).



Formalin-fixed, paraffin-embedded human Lung SqCC stained with PD-L1 Recombinant Rabbit Monoclonal Antibody (PDL1/4451R).

Specificity & Comments

PD-L1 is a checkpoint regulator in immune cells, it is expressed on immune or non-hematopoietic cells. Expression of the protein is seen during pregnancy where it has a role in suppressing the immune system. PD-L1 induces an inhibitory signal in activated T-cells and promotes T-cell apoptosis. It is overexpressed in a number of different cancers where it is believed to play a significant role in the cancer's ability to evade the immune system.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology, Mast Cell Marker, PD-1 blockade immunotherapy, Signal Transduction

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.