

Recombinant CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) Antibody

Rabbit Monoclonal Antibody [Clone C209/2749R]

Catalog No	Format	Size
30835-RBM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
30835-RBM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
30835-RBM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	C209/2749R
Gene Name	CD209
Immunogen	Recombinant full-length human CD209 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	45kDa
Cellular Localization	Cell membrane, Secreted
Species Reactivity	Human
Positive Control	THP-1 or HeLa cells (IF/FACS). Human liver or spleen lysate (WB). Human small intestine (IHC), Intestine, Liver, Lung, Heart.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant CD209 / DC-SIGN (Pathogen Receptor on Dendritic Cells) Antibody




Western blot analysis of Human Intestine, Human Heart, Human Kidney, Human Liver, and Human Skin tissue lysates using DC-SIGN Recombinant Rabbit Monoclonal Antibody (C209/2749R).



Western blot analysis of Intestine, Liver and Lung tissue lysates of different species using DC-SIGN Recombinant Rabbit Monoclonal Antibody (C209/2749R).



SDS-PAGE Analysis of Purified CD209 antigen Recombinant Rabbit Monoclonal Antibody (C209/2749R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Small Intestine stained with CD209 Recombinant Rabbit Monoclonal Antibody (C209/2749R).

Specificity & Comments

DC-SIGN is a transmembrane receptor that is expressed on the surface of dendritic cells and macrophages. It is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization, which allows the receptor to bind multivalent ligands with high avidity.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Dendritic Cell Marker, Hematopoietic Stem Cells, Immunology