

## HSP90AB1 (Heat Shock Protein 90) Antibody

Mouse Monoclonal Antibody [Clone HSP90AB1/3954]

Catalog No	Format	Size
3326-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3326-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3326-MSM4-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	HSP90AB1/3954
<b>Gene Name</b>	HSP90AB1
<b>Immunogen</b>	Recombinant human HSP90AB1 protein fragment (around aa581-704) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	~90kDa
<b>Cellular Localization</b>	Cell membrane, Cell surface, Cytoplasm, Dynein axonemal particle, Melanosome, Nucleus, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Human spleen, stomach or pancreas tissue. MCF-7 cells.

\*Optimal dilution for a specific application should be determined.



### Product Images for HSP90AB1 (Heat Shock Protein 90) Antibody



Western blot analysis of MCF-7 cell lysate using HSP90AB1 Mouse Monoclonal Antibody (HSP90AB1/3954).



SDS-PAGE Analysis of Purified HSP90AB1 Mouse Monoclonal Antibody (HSP90AB1/3954). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human pancreas stained with HSP90AB1 Mouse Monoclonal Antibody (HSP90AB1/3954).

Analysis of Protein Array containing more than 19,000 full-length human proteins using HSP90AB1 Mouse Monoclonal Antibody (HSP90AB1/3954). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

### Specificity & Comments

This gene encodes a member of the heat shock protein 90 family; these proteins are involved in signal transduction, protein folding and degradation and morphological evolution. This gene encodes the constitutive form of the cytosolic 90 kDa heat-shock protein and is thought to play a role in gastric apoptosis and inflammation. Alternative splicing results in multiple transcript variants. Pseudogenes have been identified on multiple chromosomes.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Developmental Biology, Immunology, Infectious Disease, Signal Transduction

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.