

Recombinant CD50 / ICAM3 Antibody

Rabbit Monoclonal Antibody [Clone ICAM3/2873R]

| Catalog No | Format | Size |
|-----------------|--|--------|
| 3385-RBM6-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3385-RBM6-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3385-RBM6-P1ABX | Purified Ab WITHOUT BSA or Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|---|
| Clone | ICAM3/2873R |
| Immunogen | Recombinant fragment (around aa1-547) of the human CD50 protein (exact sequence is proprietary) |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | 59.54kDa |
| Cellular Localization | Membrane |
| Species Reactivity | Human |
| Positive Control | HL-60 or THP-1 cells. Lymph node and tonsil. |

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant CD50 / ICAM3 Antibody



SDS-PAGE Analysis of Purified Intercellular adhesion molecule 3 Recombinant Rabbit Monoclonal Antibody (ICAM3/2873R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Tonsil stained with CD50 Rabbit Recombinant Monoclonal Antibody (ICAM3/2873R).

Specificity & Comments

Recognizes the D1 domain of an N-glycosylated glycoprotein of 120kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (Leucocyte Workshop VI). CD50 is the major ligand for LFA-1 (CD11a/CD18) and may have signaling role to increase adhesion. It is expressed on thymocytes and T lymphocytes and is resistant to treatment with phosphatidylinositol (PI) phospholipase C. This MAAb is excellent for staining of formalin/paraffin tissues.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

2 Union Square, Union City, CA 94587, Tel: 1-866-2-NEOBIO (1-866-263-6246)
orders@NeoBiotechnologies.com, www.NeoBiotechnologies.com