

Interferon alpha-1 (IFNA1) Antibody

Mouse Monoclonal Antibody [Clone 2-52]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 3439-MSM2-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3439-MSM2-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3439-MSM2-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|-------------------------|---------------------|------|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |

Product Details

| | |
|------------------------|---|
| Clone | 2-52 |
| Gene Name | IFNA1, IFNA13 |
| Immunogen | Purified recombinant human IFN-alpha1 |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 16-27kDa |
| Cellular Localization | Secreted |
| Species Reactivity | Human |
| Positive Control | Stimulated human peripheral blood mononuclear cells |

**Optimal dilution for a specific application should be determined.*

Product Images for Interferon alpha-1 (IFNA1) Antibody



SDS-PAGE Analysis of Purified IFNA1 Mouse Monoclonal Antibody (2-52).
Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

This MAb is specific for human interferon alpha 1 and does not cross react with human interferon alpha 2. Interferons are widely used therapeutic agents because of their anti-tumor and anti-viral effects and because of their modulatory effects on the immune system. These cytokines produce their effects by binding to the Type 1 Interferon- & Receptor (IFNAR1). Down regulation of this receptor plays a key role in determining the magnitude and duration of cytokine signaling. This down regulation is influenced by phosphorylation of Serine 535 and 539 in the IFNAR1.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Cytokine Signaling, Infectious Disease

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.