Recombinant Interleukin-1beta (IL-1beta) Antibody

Rabbit Monoclonal Antibody [Clone IL1B/7049R]

Catalog No	Format	Size
3553-RBM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3553-RBM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3553-RBM13-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	IL1B/7049R
Gene Name	IL1B
Immunogen	Recombinant fragment of human IL1B protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	31kDa (proprotein); 17kDa (mature)
Cellular Localization	Cytoplasm, Cytosol, Extracellular exosome, Lysosome, Secreted
Species Reactivity	Human
Positive Control	LPS stimulated human peripheral blood mononuclear cells. Human colon or liver tissue.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Interleukin-1beta (IL-1beta) Antibody



SDS-PAGE Analysis of Purified IL1B Recombinant Rabbit Monoclonal Antibody (IL1B/7049R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Recognizes a protein of 17-31kDa, identified as interleukin 1beta. It shows no cross-reaction with interleukin 1alpha. Epitopes of MAb IL1B/463 and IL1B/7049R are different and provide a good pair for developing an ELISA. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Cardiovascular, Cytokine Signaling, Immunology, Infectious Disease, MAPK Signaling, Neuroinflammation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.