

## IL3RA / CD123 (Acute Myeloid Leukemia Marker) Antibody

Mouse Monoclonal Antibody [Clone IL3RA/1822]


Catalog No	Format	Size
3563-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3563-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3563-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Western Blot (WB)	2-4ug/ml	

Product Details	
<b>Clone</b>	IL3RA/1822
<b>Gene Name</b>	IL3RA
<b>Immunogen</b>	Recombinant fragment of human IL3RA protein (around aa 26-171) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	70kDa
<b>Cellular Localization</b>	Membrane
<b>Species Reactivity</b>	Hamster, Human, Mouse, Rat
<b>Positive Control</b>	Human tonsil, lymph node or stomach. THP-1, RPMI-8226 or HDLM-2 cells.

\*Optimal dilution for a specific application should be determined.

### Product Images for IL3RA / CD123 (Acute Myeloid Leukemia Marker) Antibody



Western blot analysis of Human Skin, Mouse Skin, Rat Skin, Hamster Skin and Guinea pig Skin tissue lysates using IL-3Ra Mouse Monoclonal Antibody (IL3RA/1822).

### Specificity & Comments

CD123 is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudo-autosomal region on chromosomes X or Y.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

AKT Signaling, Cytokine Signaling, Hematopoietic Stem Cells, Immunology, Signal Transduction

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.