

CD137 / 4-1BB / TNFRSF9 Antibody

Mouse Monoclonal Antibody [Clone 4-1BB/3201]


Catalog No	Format	Size
3604-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3604-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3604-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


Product Details	
Clone	4-1BB/3201
Gene Name	TNFRSF9
Immunogen	A recombinant fragment (around aa 19-188) of human CD137 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	32kDa (monomer); 85kDa (dimer)
Cellular Localization	Cell surface, Membrane
Species Reactivity	Human
Positive Control	Human kidney cancer

*Optimal dilution for a specific application should be determined.



Product Images for CD137 / 4-1BB / TNFRSF9 Antibody



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Biotin-conjugated CD137 Mouse Monoclonal Antibody (4-1BB/3201).



Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Biotin-conjugated CD137 Mouse Monoclonal Antibody (4-1BB/3201).



SDS-PAGE Analysis of Purified CD137 Mouse Monoclonal Antibody (4-1BB/3201). Confirmation of Purity and Integrity of Antibody.

Analysis of Protein Array containing more than 19,000 full-length human proteins using CD137-Monospecific Mouse Monoclonal Antibody (4-1BB/3201). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

CD137 belongs to the tumor necrosis factor receptor family and delivers a costimulatory signal to T lymphocytes. It is expressed on activated T cells and binds an inducible ligand that is found on B cells, macrophages and dendritic cells. Interactions between CD137 and its ligand are involved in antigen presentation and the generation of cytotoxic T cells. CD137 antibody may improve cancer treatment, and has been implicated in breast cancer, melanoma and lymphoma.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, B Cell Markers, Cardiovascular, Cytokine Signaling, Immunology

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.