

Recombinant Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rINHA/6919]

Catalog No	Format	Size
3623-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3623-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3623-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	rINHA/6919
Gene Name	INHA
Immunogen	Recombinant fragment of human Inhibin alpha protein (around aa 233-362) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	47kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Human prostate or testis.



*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody



IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using rINHA/6919 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH 9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.

SDS-PAGE Analysis Purified Inhibin, alpha Recombinant Mouse Monoclonal (rINHA/6919). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of human testis tissue lysate using INHA Recombinant Mouse Monoclonal Antibody (INHA/6919).

IHC analysis of formalin-fixed, paraffin-embedded humantesticular carcinoma. Staining with INHA/6919at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

It recognizes a 47kDa protein, which is identified as alpha sub-unit of Inhibin. It is a gonadal protein that preferentially suppresses the secretion of pituitary follicle-stimulating hormone (FSH). Inhibin comprises two subunits, Inhibin A and Inhibin B. Each subunit consists of the same α subunit, covalently linked to 1 of 2 distinct subunits, $\beta\alpha$ or $\beta\beta$. Originally isolated from ovarian follicular fluid and characterized as a disulphide-linked dimeric glycoprotein, inhibin belongs to the transforming growth factor β (TGF β) superfamily. Antibodies against Inhibin are useful in making a differentiation between adrenal cortical tumors and renal cell carcinoma. Sex cord stromal tumors of the ovary as well as trophoblastic tumors also demonstrate cytoplasmic positivity. Inhibin antibody is also used to make the differential diagnosis of intrauterine vs. ectopic pregnancy in endometrial curetting.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular