

IRF3 Antibody

Mouse Monoclonal Antibody [Clone PCR-P-IRF3-6C8]

Catalog No	Format	Size
3661-MSM15-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3661-MSM15-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3661-MSM15-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	PCR-P-IRF3-6C8
Gene Name	IRF3
Immunogen	Recombinant full-length human IRF3 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	~50kDa
Cellular Localization	Cytoplasm, Mitochondrion, Nucleus
Species Reactivity	Human
Positive Control	HeLa, U87 or Ramos cells.

*Optimal dilution for a specific application should be determined.

Product Images for IRF3 Antibody



Flow Cytometric Analysis of PFA-fixed U87 cells. IRF3 Mouse Monoclonal Antibody (PCR-P-IRF3-6C8) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

Specificity & Comments

Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon- α and β) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon- α and interferon- β genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth; overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3 β p48 and ~~IFN consensus sequence binding protein (ICSBP).~~

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Cytokine Signaling, Infectious Disease

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
