

Arginase 1 (Hepatocellular Carcinoma Marker) Antibody

Mouse Monoclonal Antibody [Clone ARG1/1125+ ARG1/1126]


Catalog No	Format	Size
383-MSM3-P0	Purified Ab with BSA and Azide	200ug/ml
383-MSM3-P1	Purified Ab with BSA and Azide	200ug/ml
383-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml


Product Details	
Clone	ARG1/1125+ ARG1/1126
Gene Name	ARG1
Immunogen	Recombinant human ARG1 protein fragment (around aa11-97) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG3 / Kappa
Mol. Weight of Antigen	35-38kDa
Cellular Localization	Cytoplasm, Cytoplasmic granule
Species Reactivity	Human
Positive Control	293T cells. Hepatocellular Carcinoma (HCC).

**Optimal dilution for a specific application should be determined.*

Product Images for Arginase 1 (Hepatocellular Carcinoma Marker) Antibody



Formalin-paraffin human Hepatocellular Carcinoma stained with ARG1 Mouse Monoclonal Antibody (ARG1/1125 + ARG1/1126).



Formalin-paraffin human Hepatocellular Carcinoma stained with ARG1 Mouse Monoclonal Antibody (ARG1/1125 + ARG1/1126).

Specificity & Comments

Recognizes a protein of 35-38kDa, which is identified as Arginase 1 (ARG1). Arginase is a manganese metallo-enzyme that catalyzes the hydrolysis of arginine to generate ornithine and urea. Arginase I and II are isoenzymes, which differ in subcellular localization, regulation, and possibly function. Arginase I is a cytosolic enzyme, which is expressed mainly in the liver as part of the urea cycle, whereas arginase II is a mitochondrial protein found in a variety of tissues. Antibody to ARG-1 labels hepatocytes in normal tissues and granulocytes in peripheral blood. ARG-1 is a sensitive and specific marker for identification of hepatocellular carcinoma.

Research Areas

Cardiovascular, Immunology, Dendritic Cell Marker

Known Applications & Suggested Dilutions

Immunohistology (Formalin-fixed) (2-4ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
