

Cytokeratin 6 (KRT6) (Basal Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone KRT6/1702]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 3853-MSM2-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3853-MSM2-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3853-MSM2-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |



| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details

| | |
|------------------------|-------------------------------------|
| Clone | KRT6/1702 |
| Gene Name | KRT6A |
| Immunogen | Recombinant human keratin 6 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2a / Kappa |
| Mol. Weight of Antigen | 56kDa |
| Species Reactivity | Human, Mouse |
| Positive Control | Tonsil or basal cell carcinoma. |

*Optimal dilution for a specific application should be determined.

Product Images for Cytokeratin 6 (KRT6) (Basal Cell Marker) Antibody



SDS-PAGE Analysis of Purified KRT6 Mouse Monoclonal Antibody (KRT6/1702). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human Bladder Carcinoma stained with Cytokeratin 6 Monoclonal Antibody (KRT6/1702)

Specificity & Comments

This MAb recognizes a protein of 56kDa, identified as cytokeratin 6 (CK6). In humans, multiple isoforms of Cytokeratin 6 (6A-6F), encoded by several highly homologous genes, have distinct tissue expression patterns, and Cytokeratin 6A is the dominant form in epithelial tissue. The gene encoding human Cytokeratin 6A maps to chromosome 12q13, and mutations in this gene are linked to several inheritable hair and skin pathologies. Keratins 6 and 16 are expressed in keratinocytes, which are undergoing rapid turnover in the suprabasal region (also known as hyper-proliferation-related keratins). Keratin 6 is found in hair follicles, suprabasal cells of a variety of internal stratified epithelia, in epidermis, in both normal and hyper-proliferative situations. Epidermal injury results in activation of keratinocytes, which express CK6 and CK16. CK6 is strongly expressed in about 75% of head and neck squamous cell carcinomas. Expression of CK6 is particularly associated with differentiation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.