

Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone KRT7/1198]

Catalog No	Format	Size
3855-MSM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3855-MSM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3855-MSM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	KRT7/1198
Gene Name	KRT7
Immunogen	Recombinant full-length human KRT7 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	55kDa
Cellular Localization	Cytoplasm
Species Reactivity	Human, Rat
Positive Control	cervix or breast., HeLa cells. Carcinoma of ovary, Lung


*Optimal dilution for a specific application should be determined.

Product Images for Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody




Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).


Western Blot Analysis of human HeLa cell lysate using Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).




Formalin-fixed, paraffin-embedded Rat Lung stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).



Immunofluorescence Analysis of MeOH-fixed HeLa cells labeling KRT7 with KRT7 Mouse Monoclonal Antibody (KRT7/1198) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Redot.



SDS-PAGE Analysis of Purified KRT7 Mouse Monoclonal Antibody (KRT7/1198). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Lung Carcinoma stained with Cytokeratin 7 Mouse Monoclonal Antibody (KRT7/1198).

Specificity & Comments

It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This MAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or gastrointestinal tract. This MAb is highly useful in distinguishing ovarian carcinomas (keratin 7+) from colon carcinomas (keratin 7-).

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Autophagy, Developmental Biology

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.