

Recombinant Cytokeratin 8 (KRT8) Antibody

Mouse Monoclonal Antibody [Clone rKRT8/6471]


Catalog No	Format	Size
3856-MSM21-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3856-MSM21-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3856-MSM21-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	


Product Details	
Clone	rKRT8/6471
Immunogen	Cytoskeleton preparation containing cytokeratin 8
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	53.7kDa
Cellular Localization	Cytoplasm, Nucleoplasm, Nucleus, Nucleus matrix
Species Reactivity	Human
Positive Control	Colon, HCT116, lung or breast carcinoma., MCF-7 or A431 cells. Human skin, Pancreas.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant Cytokeratin 8 (KRT8) Antibody




Formalin-fixed, paraffin-embedded human colon stained with Cytokeratin 8 Recombinant Mouse Monoclonal Antibody (rKRT8/6471).




IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using rKRT8/6471 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.




SDS-PAGE Analysis of Purified CK8 Recombinant Mouse Monoclonal Antibody (rKRT8/6471). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Pancreas and MCF7 lysates using Cytokeratin 8 Recombinant Mouse Monoclonal Antibody (rKRT8/6471).



Western blot analysis of human pancreas tissue lysates using KRT8 Recombinant Mouse Monoclonal Antibody (rKRT8/6471).



Formalin-fixed, paraffin-embedded human colon stained with Cytokeratin 8 Recombinant Mouse Monoclonal Antibody (rKRT8/6471).

Specificity & Comments

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenocarcinomas and ductal carcinomas. It is absent in squamous cell carcinomas. Hepatocellular carcinomas are defined by the use of antibodies that recognize only cytokeratin 8 and 18. CK8 exists on several types of normal and neoplastic epithelia, including many ductal and glandular epithelia such as colon, stomach, small intestine, trachea, and esophagus as well as in transitional epithelium. Anti-CK8 does not react with skeletal muscle or nerve cells. Epithelioid sarcoma, chordoma, and adamantinoma show strong positivity corresponding to that of simple epithelia (with antibodies against CK8, CK18 and CK19). Anti-CK8 is useful for the differentiation of lobular (ring-like, perinuclear) from ductal (peripheral-predominant) carcinoma of the breast.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.