

## Recombinant Cytokeratin 18 (KRT18) Antibody

Rabbit Monoclonal Antibody [Clone KRT18/2808R]

Catalog No	Format	Size
3875-RBM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3875-RBM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3875-RBM11-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	KRT18/2808R
<b>Gene Name</b>	KRT18
<b>Immunogen</b>	Recombinant human full-length KRT18 protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	45kDa
<b>Cellular Localization</b>	Cytoplasm, Nucleolus, Nucleus, Perinuclear region
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	A431 cells. Breast Cancer., HCT116, HeLa, MCF-7

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant Cytokeratin 18 (KRT18) Antibody




Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R).




Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R).




SDS-PAGE Analysis Purified CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R). Confirmation of Purity and Integrity of Antibody.




Western Blot Analysis of human HCT116 cell lysate using CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R).



Immunofluorescence Analysis of MCF-7 cells labeling CK18 with CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R) followed by Goat anti-Mouse IgG-CF488 (Green). The nuclear counterstain is Reddot (Red).



Flow Cytometric Analysis of HeLa cells using CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R) followed by Goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).



Formalin-fixed, paraffin-embedded human Thyroid Carcinoma stained with CK18 Rabbit Recombinant Monoclonal Antibody (KRT18/2808R).

### Specificity & Comments

This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs to the type A (acidic) subfamily of low molecular weight keratins, exists in combination with cytokeratin 8. It is reported that tissues from gastrointestinal tract are positive for both cytokeratin 8 and 18 but do not contain cytokeratin 14. Tissues from gastrointestinal tract, respiratory tract and urogenital tract, as well as endocrine and exocrine tissues and mesothelial cells are positive for cytokeratin 18.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Autophagy, Developmental Biology