

Lysozyme (Histiocytoma & Monocytic Acute Leukemia Marker) Antibody

Mouse Monoclonal Antibody [Clone LYZ/3942]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 4069-MSM2-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 4069-MSM2-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 4069-MSM2-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|---|
| Clone | LYZ/3942 |
| Gene Name | LYZ |
| Immunogen | Recombinant fragment of human Lysozyme protein (around aa 18-147) (exact sequence is proprietary) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2b / Kappa |
| Mol. Weight of Antigen | 17kDa |
| Cellular Localization | Secreted |
| Species Reactivity | Human |
| Positive Control | Human tonsil, lung or kidney., Spleen |

*Optimal dilution for a specific application should be determined.


Product Images for Lysozyme (Histiocytoma & Monocytic Acute Leukemia Marker) Antibody




Formalin-fixed, paraffin-embedded human tonsil stained with Lysozyme Mouse Monoclonal Antibody (LYZ/3942).




SDS-PAGE Analysis Purified Lysozyme Mouse Monoclonal Antibody (LYZ/3942). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of human spleen tissue and HepG2 cell lysates using Lysozyme Mouse Monoclonal Antibody (LYZ/3942).



Analysis of Protein Array containing more than 19,000 full-length human proteins using Lysozyme-Monospecific Mouse Monoclonal Antibody (LYZ/3942). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human spleen stained with Lysozyme Mouse Monoclonal Antibody (LYZ/3942). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Lysozyme is an enzyme, commonly referred to as the body's own antibiotic since it kills bacteria. Natural substrate of lysozyme is the bacterial cell wall peptidoglycan (cleaving the beta[1-4]glycosidic linkages between N-acetylmuramic acid and N-acetylglucosamine). Lysozyme is one of the antimicrobial agents found in human milk, and is also present in spleen, lung, kidney, white blood cells, plasma, saliva, and tears. The protein has antibacterial activity against a number of bacterial species. Lysozyme is synthesized predominantly in reactive histiocytes rather than in resting, unstimulated phagocytes. This antibody labels myeloid cells, histiocytes, granulocytes, macrophages and monocytes. It is helpful in the identification of myeloid or monocytic nature of acute leukemia.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Dendritic Cell Marker, Immunology