

Recombinant TACSTD2 / TROP2 (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone rTACSTD2/6395]


Catalog No	Format	Size
4070-MSM5-P0	Purified Ab with BSA and Azide	200ug/ml
4070-MSM5-P1	Purified Ab with BSA and Azide	200ug/ml
4070-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml


Product Details	
Clone	rTACSTD2/6395
Gene Name	TACSTD2
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	40kDa
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	HT29 cells. Human breast or colon carcinoma.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant TACSTD2 / TROP2 (Epithelial Marker) Antibody




Formalin-fixed, paraffin-embedded human liver in colon carcinoma stained with TACSTD2 Recombinant Mouse Monoclonal Antibody (rTACSTD2/6395).



IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using rTACSTD2/6395 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified TACSTD2 Recombinant Mouse Monoclonal (rTACSTD2/6395). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human colon carcinoma stained with TACSTD2 Recombinant Mouse Monoclonal Antibody (rTACSTD2/6395).

Specificity & Comments

TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites near the C-terminus. It plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells, cornea and multi-stratified epithelia. It is also highly expressed in several types of tumors and is involved in regulating the growth of carcinoma cells.

Known Applications & Suggested Dilutions

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.