

Recombinant Ep-CAM / CD326 (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone rEGP40/1110]

Catalog No	Format	Size
4072-MSM23-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4072-MSM23-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4072-MSM23-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	rEGP40/1110
Gene Name	EPCAM
Immunogen	Recombinant human EpCAM fragment from the cytoplasmic domain (around aa 289-314) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	40-43kDa
Cellular Localization	Cell junction, Lateral cell membrane, Tight junction
Species Reactivity	Human, Mouse, Rat
Positive Control	HT29 cells. Human colon or ovariancarcinoma.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant Ep-CAM / CD326 (Epithelial Marker) Antibody



Analysis of Protein Array containing more than 19,000 full-length human proteins using EpCAM Monospecific Recombinant Mouse Monoclonal Antibody (rEGP40/1110). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



SDS-PAGE Analysis of Purified Epithelial cell adhesion molecule Recombinant Mouse Monoclonal Antibody (rEGP40/1110). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human ovarian carcinoma stained with EpCAM Mouse Recombinant Monoclonal Antibody (rEGP40/1110).

Specificity & Comments

EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Stem Cell Differentiation
