

## Recombinant EpCAM / CD326 (Epithelial Marker) Antibody

Rabbit Monoclonal Antibody [Clone EGP40/2571R]

Catalog No	Format	Size
4072-RBM24-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4072-RBM24-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4072-RBM24-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	EGP40/2571R
<b>Immunogen</b>	Recombinant full-length rat EpCAM protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	34.93kDa
<b>Cellular Localization</b>	Cell junction, Lateral cell membrane, Tight junction
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	MCF-7 or HT29 cells. Breast tumor.

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant EpCAM / CD326 (Epithelial Marker) Antibody




Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with EpCAM Rabbit Recombinant Monoclonal Antibody (EGP40/2571R).



Flow Cytometric Analysis of MCF-7 cells using EpCAM Rabbit Recombinant Monoclonal Antibody (EGP40/2571R) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



SDS-PAGE Analysis of Purified Epithelial cell adhesion molecule Recombinant Rabbit Monoclonal Antibody (EGP40/2571R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with EpCAM Rabbit Recombinant Monoclonal Antibody (EGP40/2571R).

### Specificity & Comments

Epitope of this MAb is mapped between aa 202-212 of EGP40, which is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. Antibody to Ep-CAM has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.