Recombinant MCM7 (Proliferation Marker) Antibody

Mouse Monoclonal Antibody [Clone rMCM7/1468]

Catalog No	Format	Size
4176-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4176-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4176-MSM9-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rMCM7/1468
Immunogen	Recombinant fragment (around aa195-319) of the human MCM7 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	81.31kDa
Cellular Localization	Chromosome, Nucleus
Species Reactivity	Human
Positive Control	A-431, A549, HEK-293, HeP2, MCF-7, Raji or Ramos cells. Tonsil (IHC)., SKBR3, Skin.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant MCM7 (Proliferation Marker) Antibody




SDS-PAGE Analysis Purified MCM7 Recombinant Mouse Monoclonal Antibody (rMCM7/1468). Confirmation of Integrity and Purity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteinsusing MCM7 Recombinant Mouse Monoclonal Antibody (rMCM7/1468) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Western Blot Analysis of Human Skin and A549 lysates using MCM7 Mouse Recombinant Monoclonal Antibody (rMCM7/1468).



Formalin-fixed, paraffin-embedded human Tonsil stained with MCM7 Recombinant Mouse Monoclonal Antibody (rMCM7/1468).

Specificity & Comments

The specificity of this monoclonal antibody to its intended target was validated by HuProtTM Array, containing more than 19,000, full-length human proteins. MCM7 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that is essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumor suppressor protein RB1/RB.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.