

## Recombinant Ki-67 (Proliferating Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rMKI67/6615]

Catalog No	Format	Size
4288-MSM25-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4288-MSM25-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4288-MSM25-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	rMKI67/6615
<b>Immunogen</b>	Recombinant fragment (around aa2293-2478) of the human Ki67 protein (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	358.69kDa
<b>Cellular Localization</b>	Chromosome, Nucleolus, Nucleus
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Any actively proliferating cells. Human skin, tonsil or lymph node. Raji.

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant Ki-67 (Proliferating Cell Marker) Antibody




Formalin-fixed, paraffin-embedded human lymph node stained with Ki67 Recombinant Mouse Monoclonal Antibody (rMKI67/6615).




IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using rMKI67/6615 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified Ki67 Recombinant Mouse Monoclonal Antibody (rMKI67/6615). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Raji lysate using Ki67 Recombinant Mouse Monoclonal Antibody (rMKI67/6615).



Formalin-fixed, paraffin-embedded human tonsil stained with Ki67 Recombinant Mouse Monoclonal Antibody (rMKI67/6615).

### Specificity & Comments

Ki-67 antigen is a nuclear, non-histone protein that is present in all stages of the cell cycle except G0. This characteristic makes Ki-67 an excellent marker for proliferating cells and is commonly used as one of the prognostic factors in cancer studies. A correlation has been demonstrated between Ki-67 index and the histo-pathological grade of neoplasms. Assessment of Ki-67 expression in renal and ureter tumors shows a correlation between tumor proliferation and disease progression, thus making it possible to differentiate high-risk patients. Ki-67 expression may also prove to be important for distinguishing between malignant and benign peripheral nerve sheath tumors. Ki-67 labeling index has been shown to be a prognostic marker in a number of neoplasms including grade II astrocytoma, oligodendroglioma, colon carcinoma, and breast carcinoma. In general, Ki-67 is a good marker of proliferating cell populations.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.