

## Recombinant MUC6 (Mucin 6 / Gastric Mucin) Antibody

Rabbit Monoclonal Antibody [Clone MUC6/1553R]

Catalog No	Format	Size
4588-RBM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4588-RBM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4588-RBM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


Clone	MUC6/1553R
Gene Name	MUC6
Immunogen	Recombinant full-length human MUC6 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	252kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Human stomach tissue (IHC).

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant MUC6 (Mucin 6 / Gastric Mucin) Antibody



Formalin-fixed, paraffin-embedded human Pancreas stained with MUC6 Rabbit Recombinant Monoclonal Antibody (MUC6/1553R).



SDS-PAGE Analysis of Purified MUC6 Rabbit Recombinant Monoclonal Antibody (MUC6/1553R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Gastric Carcinoma stained with MUC6 Rabbit Recombinant Monoclonal Antibody (MUC6/1553R).

---

### Specificity & Comments

The MUC6 gastric mucin is a secreted glycoprotein that plays an essential role in epithelial cyto-protection from acid, proteases, pathogenic microorganisms, and mechanical trauma in the gastrointestinal tract. Mucin 6 expression is highest in the stomach and gall bladder, with lower expression in the terminal ileum and right colon. In gastric cancer, Mucin 6 has an altered expression. In normal stomach, Mucin 6 is associated with Lewis type 2; Mucin 6 is also expressed in gastric metaplasia, duodenum and pancreas. Mucin 6 is a secretory mucin, located in the deeper mucosal folds of human gall bladder, and its expression is altered with increasing degrees of inflammation.

---

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### Supplied As

200ug/ml of Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### Research Areas

Immunology, Infectious Disease

---