

## c-Myc Oncoprotein Antibody

Mouse Monoclonal Antibody [Clone 9E10.3]

Catalog No	Format	Size
4609-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4609-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4609-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	9E10.3
<b>Gene Name</b>	MYC
<b>Immunogen</b>	A synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNSCA) from C-terminus of human c-myc, coupled to KLH.
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	62-64kDa
<b>Cellular Localization</b>	Nucleolus, Nucleoplasm, Nucleus
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	MCF7 cells. Cervical Carcinoma.

\*Optimal dilution for a specific application should be determined.


### Product Images for c-Myc Oncoprotein Antibody



SDS-PAGE Analysis of Purified c-Myc Mouse Monoclonal Antibody (9E10.3). Confirmation of Integrity and Purity of Antibody



Western blot analysis of MCF7 cell lysate using c-Myc Mouse Monoclonal Antibody (9E10.3).



Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with c-Myc Mouse Monoclonal Antibody (9E10.3).

---

### Specificity & Comments

It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease.

---

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### Research Areas

Apoptosis, Autophagy, Breast Cancer, Cancer, Cardiovascular, Developmental Biology, Hypoxia, Immunology, Bladder Cancer, Colon Cancer, Cytokine Signaling, Infectious Disease, Lung Cancer, MAPK Signaling, Nuclear Marker, Signal Transduction, Stem Cell Differentiation, Transcription Factors

---

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---