

CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone NCAM1/795]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 4684-MSM5-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 4684-MSM5-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 4684-MSM5-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|--------------------------------------|
| Clone | NCAM1/795 |
| Gene Name | NCAM1 |
| Immunogen | Recombinant human NCAM1 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 145 and 125kDa 180 |
| Cellular Localization | Cell membrane, Secreted |
| Species Reactivity | Human, Rat |
| Positive Control | Cerebellum, Neuroblastoma., Pancreas |

*Optimal dilution for a specific application should be determined.


Product Images for CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Antibody




Formalin-fixed, paraffin-embedded human Lung Carcinoma stained with CD56 Mouse Monoclonal Antibody (NCAM1/795).



Western Blot Analysis of HCT116 cell lysate using CD56 Mouse Monoclonal Antibody (NCAM1/795).



SDS-PAGE Analysis of Purified CD56 Mouse Monoclonal Antibody (NCAM1/795). Confirmation of Integrity and Purity of Antibody



Formalin-fixed, paraffin-embedded human Pancreas stained with CD56 Mouse Monoclonal Antibody (NCAM1/795).

Specificity & Comments

This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Autophagy, Cardiovascular, Cytokine Signaling, Developmental Biology, Hematopoietic Stem Cells, Immunology, Mesenchymal Stem Cell Differentiation, Neural Stem Cells, Neuroscience, Signal Transduction, Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.