

Recombinant NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody

Mouse Monoclonal Antibody [Clone rNGFR/1965]

Catalog No	Format	Size
4804-MSM10-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4804-MSM10-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4804-MSM10-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rNGFR/1965
Gene Name	NGFR
Immunogen	Recombinant human p75 NGFR protein fragment (around aa 281-421) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	75kDa
Cellular Localization	Cell membrane, Cell projection, Dendritic spine, Growth cone, Perikaryon
Species Reactivity	Human
Positive Control	Neurofibroma. Soma and axons of sensory neurons and ganglionic satellite cells. Melanomas., Neuronal axons, Schwann cells and perineural cells of peripheral nerves or tumors of nerve sheath differentiation e. G. Schwannoma

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody




Formalin-fixed, paraffin-embedded human melanoma stained with NGFR Recombinant Mouse Monoclonal Antibody (rNGFR/1965).



Western Blot Analysis of U87 cell lysate using NGFR Recombinant Mouse Monoclonal Antibody (rNGFR/1965).



SDS-PAGE Analysis of Purified Tumor necrosis factor receptor superfamily member 16 Recombinant Mouse Monoclonal Antibody (rNGFR/1965). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human melanoma stained with NGFR Recombinant Mouse Monoclonal Antibody (rNGFR/1965).

Specificity & Comments

It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Mesenchymal Stem Cell Differentiation, Neuroscience, Nuclear Marker, Signal Transduction, Stem Cell Differentiation