

## NRF1 Antibody

Mouse Monoclonal Antibody [Clone NRF1/2609]

Catalog No	Format	Size
4899-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4899-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4899-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details

<b>Clone</b>	NRF1/2609
<b>Gene Name</b>	NRF1
<b>Immunogen</b>	Recombinant full-length human NRF1 protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	30kDa (bZIP region); 65-120kDa (glycosylated)
<b>Cellular Localization</b>	Nucleus
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Ubiquitous expression, strongest in skeletal muscle.


*\*Optimal dilution for a specific application should be determined.*

### Product Images for NRF1 Antibody




Western Blot Analysis of human HEK293, HeLa, A431, MOLT4 cell lysates using NRF1 Mouse Monoclonal Antibody (NRF1/2609).

Formalin-fixed, paraffin-embedded human colon carcinoma stained with NRF1 Mouse Monoclonal Antibody (NRF1/2609).



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with NRF1 Mouse Monoclonal Antibody (NRF1/2609).



SDS-PAGE Analysis of Purified NRF1 Mouse Monoclonal Antibody (NRF1/2609). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

The NF-E2 DNA binding protein is composed of two subunits, p45 and MafK, and it regulates expression of globin genes in developing erythroid cells through interaction with Maf recognition elements (MAREs). A family of NF-E2 related proteins, which are collectively known as the Cap 'n' collar (CNC) family and include Nrf1 (also designated TCF11), Nrf2 and Nrf3, are bZIP transcription factors that heterodimerize with Maf proteins to bind MARE sequences. The Nrf proteins also bind the antioxidant response element (ARE) and are implicated in the regulation of detoxification enzymes and the oxidative stress response. They do so by heterodimerizing with Jun family members (c-Jun, JunB and JunD) to activate gene expression, specifically the detoxifying enzyme, NQO1. Nrf2 is widely expressed and is thought to translocate to the nucleus after treatment with xenobiotics and antioxidants, which stimulate its release from a repressor protein Keap1. Nrf3 is highly expressed in placenta, B cells and monocytes.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Nuclear Marker