PAX7 (Rhabdomyosarcoma Marker) Antibody

Mouse Monoclonal Antibody [Clone PAX7/1187]

Catalog No	Format	Size
5081-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug
5081-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5081-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	PAX7/1187
Gene Name	PAX7
Immunogen	Recombinant fragment (aa301-505) of human PAX7 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	57kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Human rhabdomyosarcoma (RMS)or testicular carcinoma.

**Optimal dilution for a specific application should be determined.*


Product Images for PAX7 (Rhabdomyosarcoma Marker) Antibody




Formalin-fixed, paraffin-embeddedhuman testicular carcinoma stained with PAX7 Mouse Monoclonal Antibody (PAX7/1187).



Formalin-fixed, paraffin-embeddedhuman testicular carcinoma stained with PAX7 Mouse Monoclonal Antibody (PAX7/1187).



SDS-PAGE Analysis of Purified Paired box protein Pax-7 Mouse Monoclonal Antibody (PAX7/1187). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embeddedhuman rhabdomyosarcoma stained with PAX7 Mouse Monoclonal Antibody(PAX7/1187).

Specificity & Comments

The Pax gene family of nuclear transcription factors is comprised of nine members that function during embryogenesis to regulate the temporal and position-dependent differentiation of cells. In addition, the family is involved in a variety of signal transduction pathways in the adult organism. Mutations in the Pax family of proteins have been linked to disease and cancer in humans. Pax-7 is a protein specifically expressed in cultured satellite cell-derived myoblasts. In situ hybridization reveals that Pax-7 is also expressed in satellite cells residing in adult muscle. A chromosomal aberration in the gene encoding Pax-7 causes rhabdomyosarcoma 2 (RMS2) (also called alveolar rhabdomyosarcoma).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Mesenchymal Stem Cell Differentiation

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.