

PCNA (Proliferating Cell Nuclear Antigen) (G1- & S-phase Marker) Antibody

Mouse Monoclonal Antibody [Clone PC10]

Catalog No	Format	Size
5111-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5111-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5111-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	PC10
Gene Name	PCNA
Immunogen	Rat PCNA/protein A fusion protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	36kDa
Cellular Localization	Nucleus
Species Reactivity	Chicken, Drosophila melanogaster, Guinea Pig, Hamster, Human, Monkey, Mouse, Pig, Pombe, Rabbit, Rat, Yeast (S. pombe and S. cerevisiae), Zebrafish
Positive Control	A-431, 293T, Hep G2, MCF7, NIH/3T3, A549, Raji cells, Human Tonsil, Human Spleen or kidney

**Optimal dilution for a specific application should be determined.*


Product Images for PCNA (Proliferating Cell Nuclear Antigen) (G1- & S-phase Marker) Antibody




SDS-PAGE Analysis of Purified PCNA Mouse Monoclonal Antibody (PC10).




Flow Cytometric Analysis of A549 cells using PCNA Mouse Monoclonal Antibody (PC10) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).




Western Blot Analysis of A-431, 293T, Hep G2, MCF7 and NIH/3T3 cell lysates using PCNA Mouse Monoclonal Antibody (PC10).



Western Blot Analysis of Human Kidney, Mouse Kidney, Rat Kidney, Hamster Kidney, Guinea pig Kidney and Rabbit Kidney tissue lysates using PCNA Mouse Monoclonal Antibody (PC10).



Western Blot Analysis of Human Spleen, Mouse Spleen, Rat Spleen and Ramos lysate using PCNA Mouse Monoclonal Antibody (PC10).



Formalin-fixed, paraffin-embedded human Tonsil stained with PCNA Mouse Monoclonal Antibody (PC10).

Specificity & Comments

Recognizes a non-histone protein of 36kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during late G1 phase immediately before the onset of DNA synthesis, becoming maximal during S-phase and declining during G2 and M phases. This MAb is excellent for multiple applications.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Nuclear Marker, Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.