PDCD1 / PD1 / CD279 (Programmed Cell Death 1) Antibody

Mouse Monoclonal Antibody [Clone PDCD1/7125]

Catalog No	Format	Size
5133-MSM12-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5133-MSM12-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5133-MSM12-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	PDCD1/7125
Gene Name	PDCD1
Immunogen	Recombinant fragment (around aa1-200) of human PDCD1 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	55kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	Human tonsil, lymph node or appendix.

**Optimal dilution for a specific application should be determined.*


Product Images for PDCD1 / PD1 / CD279 (Programmed Cell Death 1) Antibody




IHC analysis of FFPE human skin (RNA expression: 0.2 NX). Negative tissue control using PDCD1/7125 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis Purified PD1 Mouse Monoclonal Antibody (PDCD1/7125). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Tonsil and Brain tissue lysate using PD1 Mouse Monoclonal Antibody (PDCD1/7125).



Formalin-fixed, paraffin-embedded human tonsil stained with PD1 (CD279) Mouse Monoclonal Antibody (PDCD1/7125). HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

PDCD-1 (programmed cell death-1 protein), also designated CD279, is a type I transmembrane receptor and a member of the immunoglobin gene superfamily. It is expressed on activated T-cells, B-cells, and myeloid cells. Anti-PDCD-1 is a marker of angioimmunoblastic lymphoma and suggests a unique cell of origin for this neoplasm. Unlike CD10 and BCL6, PDCD-1 is expressed by few B-cells, so anti-PDCD-1 may be a more specific and useful diagnostic marker in angioimmunoblastic lymphoma. In addition, PDCD-1 expression provides evidence that angioimmunoblastic lymphoma is a neoplasm derived from germinal center-associated T-cells.

Supplied As

200ug/ml of Ab purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

B Cell Markers, Cardiovascular, Immunology, Infectious Disease, PD-1 blockade immunotherapy

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.