

POLE3 / CHRAC17 Antibody

Mouse Monoclonal Antibody [Clone PCRP-POLE3-2F10]

Catalog No	Format	Size
54107-MSM2-P0	Purified Ab with BSA and Azide	200ug/ml
54107-MSM2-P1	Purified Ab with BSA and Azide	200ug/ml
54107-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells
Immunofluorescence (IF)	1-3ug/ml
Immunohistochemistry (IHC)	1-2ug/ml

Product Details

Clone	PCRP-POLE3-2F10				
Gene Name	POLE3				
Immunogen	Recombinant full-length human POLE3 protein				
Host	Mouse				
Clonality	Monoclonal				
Isotype / Light Chain	lgG1				
Mol. Weight of Antigen	16.86kDa				
Cellular Localization	Nucleus				
Species Reactivity	Human				
Positive Control	HeLa or K562 cells.				

*Optimal dilution for a specific application should be determined.

Product Images for POLE3 / CHRAC17 Antibody



Formalin-fixed, paraffin-embedded human lymphoma stained with POLE3 Mouse Monoclonal Antibody (PCRP-POLE3-2F10) at 2ug/ml. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

k	Da		R	NR	
250	_	-			
150	_	hours		-	
100	_	1	-		
75	_	-			2ug loading
50	_	-	-		reduced
37	_				K=reduced
25	_		-		
20	_				
15	_	_	_		
10	_	-			

SDS-PAGE Analysis. Purified POLE3 Mouse Monoclonal Antibody (PCRP-POLE3-2F10). Confirmation of Purity and Integrity of Antibody.





Analysis of Protein Array containing more than 19,000 full-length human proteinsusing POLE3 Mouse Monoclonal Antibody (PCRP-POLE3-2F10). Z- and S- Score: The Z- score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Flow Cytometric Analysis of PFA-fixed K562 cells. POLE3 Mouse Monoclonal Antibody (PCRP-POLE3-2F10) followed by goat anti-mouse IgG-CF488 (blue); isotype control (red).

Specificity & Comments

DNA replication is initiated by the binding of initiation factors to the origin of replication. Nucleosomes inhibit access to the replication machinery at these origin sequences. Nucleosome remodeling factors increase the accessibility of nucleosomal DNA to transcriptional regulators. CHRAC15 and CHRAC17 are subunits of the nucleosomal remodeling factor CHRAC (chromatin accessibility complex), which increases the accessibility of nucleosomal DNA in an ATP-dependent manner. Unlike other known chromatin remodeling factors, CHRAC also functions during chromatin assembly by using ATP to convert irregular chromatin into a regular array of nucleosomes with even spacing. This conversion process occurs when CHRAC organizes randomly deposited histones into a regularly spaced array. In the presence of CHRAC, the nucleosomal ATPase ISWI catalyzes several ATP-dependent transitions of chromatin structure.

Research Areas

Nuclear Marker

Known Applications & Suggested Dilutions

Flow Cytometry (1-2ug/million cells) | Immunofluorescence (1-2ug/ml) | ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

