

## Recombinant OCT-2 (POU2F2) (B-Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone OCT2/7073R]



Catalog No	Format	Size
5452-RBM5-P0	Purified Ab with BSA and Azide	200ug/ml
5452-RBM5-P1	Purified Ab with BSA and Azide	200ug/ml
5452-RBM5-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml

Product Details	
<b>Clone</b>	OCT2/7073R
<b>Gene Name</b>	POU2F2
<b>Immunogen</b>	Recombinant fragment of human OCT2 protein (around aa 112-297) (exact sequence is proprietary)
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	60kDa
<b>Cellular Localization</b>	Cytoplasm, Nucleus
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Raji or U-937 cells. Human spleen or tonsil.

*\*Optimal dilution for a specific application should be determined.*

### Product Images for Recombinant OCT-2 (POU2F2) (B-Cell Marker) Antibody



Formalin-fixed, paraffin-embedded human tonsil stained with Oct-2 Recombinant Mouse Monoclonal Antibody (rOCT2/7196). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Formalin-fixed, paraffin-embedded human tonsil stained with Oct-2 Recombinant Mouse Monoclonal Antibody (rOCT2/7196). Inset: PBS instead of primary antibody; secondary only negative control.

### Specificity & Comments

Oct-2 is a transcription factor of the POU homeo-domain family that binds to the Ig gene octamer sites, regulating B-cell-specific genes. Oct-2 expression can be used as a marker of B-cell lineage and differentiation. Germinal center B-cells, mantle B-cells, monocytoid B-cells, and plasma cells show high level expression of Oct-2. Additionally, mantle cell lymphoma, follicular lymphoma, marginal zone lymphoma, plasmacytoma, Burkitt lymphoma, diffuse large cell lymphoma, diffuse large B-cell lymphoma, Hodgkin lymphoma display increased expression of Oct-2. Several studies of Oct-2 expression have shown a low level expression in pre-B, T-cell, myelomonocytic, and epithelial cell lines, whereas all mature B-cell lines have high levels of expression. In spite of scanty evidence for Oct-2 expression in T cells, it is believed that this factor participates in transcriptional regulation during T-cell activation.

---

### Research Areas

Nuclear Marker, Transcription Factors

---

### Known Applications & Suggested Dilutions

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

---

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---