Recombinant CD45RO (T-Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone PTPRC/1975R]

Catalog No	Format	Size
5788-RBM25-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5788-RBM25-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5788-RBM25-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	PTPRC/1975R
Immunogen	Isolated neoplastic cells from T cell lymphoma
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	147.48kDa
Cellular Localization	Cell membrane, Membrane raft, Synapse
Species Reactivity	Human
Positive Control	Ramos, U-698 or GA-10 cells. Tonsil and Spleen.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant CD45RO (T-Cell Marker) Antibody




Western Blot Analysis of Human spleen tissue lysate using CD45 Rabbit Recombinant Monoclonal Antibody (PTPRC/1975R).




Flow Cytometric Analysis of paraformaldehyde-fixed Raji cells using CD45 Rabbit Recombinant Monoclonal Antibody (PTPRC/1975R) followed by Goat anti-rabbit IgG-CF488 (Blue); Isotype Control (Red).



Paraformaldehyde-fixed Raji cells stained with CD45 Rabbit Recombinant Monoclonal Antibody (PTPRC/1975R) followed by Goat anti-rabbit IgG-CF488 (Green). Counterstain is Phalloidin (red).



SDS-PAGE Analysis of Purified Receptor-type tyrosine-protein phosphatase C Recombinant Rabbit Monoclonal Antibody (PTPRC/1975R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Lymph Node stained with CD45 RabbitRecombinant Monoclonal Antibody (PTPRC/1975R).

Specificity & Comments

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.