

RET Proto-oncogene Antibody

Mouse Monoclonal Antibody [Clone RET/2663]

Catalog No	Format	Size
5979-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5979-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5979-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note

Product Details

Clone	RET/2663
Gene Name	RET
Immunogen	Recombinant fragment (around aa 702-848) of human RET protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	150kDa (precursor); 170kDa (Mature)
Cellular Localization	Cell membrane, Endosome membrane
Species Reactivity	Human
Positive Control	Breast, Prostate or Colon Carcinoma.


*Optimal dilution for a specific application should be determined.

Product Images for RET Proto-oncogene Antibody




Formalin-fixed, paraffin-embedded human prostate carcinoma stained with RET Mouse Monoclonal Antibody (RET/2663).

Analysis of Protein Array containing more than 19,000 full-length human proteins using RET Mouse Monoclonal Antibody (RET/2663). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (Monoclonal Antibody) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a Monoclonal Antibody to its intended target. A Monoclonal Antibody is considered to be specific to its intended target, if the Monoclonal Antibody has an S-score of at least 2.5. For example, if a Monoclonal Antibody binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that Monoclonal Antibody to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with RET Mouse Monoclonal Antibody (RET/2663).



SDS-PAGE Analysis of Purified RET Mouse Monoclonal Antibody (RET/2663). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

The Ret proto-oncogene is structurally related to the growing family of tyrosine kinase transmembrane receptors and is involved in GDNF signaling. RET expression is reported in several regions of the central nervous system; in the developing cranial nerve ganglia and a subset of cells within dorsal root ganglia, in motor neurons in the spinal cord and hindbrain, in neuro-retina and the growing tips of the renal collecting ducts in developing kidney. Alterations in RET gene are associated with diseases including papillary thyroid carcinoma, multiple endocrine neoplasia (type 2A and 2B), familial medullary thyroid carcinoma, and a congenital developmental disorder known as Hirschsprung's disease.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Developmental Biology, Neuroscience, Signal Transduction, Transcription Factors