

Calprotectin / MRP14 / S100A9 / Calgranulin B Antibody

Mouse Monoclonal Antibody [Clone MRP14/840]

Catalog No	Format	Size
6280-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6280-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6280-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	MRP14/840
Gene Name	S100A9
Immunogen	Recombinant human MRP14 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgM / Kappa
Mol. Weight of Antigen	14kDa
Cellular Localization	Cell membrane, Cytoplasm, Cytoskeleton, Secreted
Species Reactivity	Human
Positive Control	lymph node, or spleen, Tonsil

**Optimal dilution for a specific application should be determined.*

Product Images for Calprotectin / MRP14 / S100A9 / Calgranulin B Antibody



Formalin-fixed, paraffin-embedded human tonsil stained with MRP14 Monoclonal Antibody (MRP14/840)

Specificity & Comments

Recognizes a protein of 14kDa, identified as MRP-14 (also known as Calgranulin B or S100AA9). It comprises 60% of the cytoplasmic protein fraction of circulating polymorphonuclear granulocytes and is also found in monocytes, macrophages and ileal tissue eosinophils. Peripheral blood monocytes carry the antigen extra- and intracellularly, neutrophils only intracellularly. It is a potent chemotactic factor for neutrophils. Plasma concentrations are elevated in diseases associated with increased neutrophil activity, like inflammatory bowel disease. Granulocytes terminate their existence after transmigration through the intestinal wall. Therefore, it is also detectable in feces. Elevated levels have been observed in body fluids such as plasma, saliva, gingival crevicular fluid, stools, and synovial fluid during infection and inflammatory conditions. This MAb reacts with neutrophils, monocytes, and macrophages, and has been shown as an important marker for identifying macrophages in tissue sections.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate. Prepared in 1mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Dendritic Cell Marker, Immunology, Infectious Disease, Signal Transduction
