Recombinant StAR (Steroidogenic Acute Regulator) (Leydig Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone STAR/3915R]

Catalog No	Format	Size
6770-RBM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6770-RBM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6770-RBM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	STAR/3915R
Gene Name	STAR
Immunogen	Recombinant fragment (around aa 39-108) of human STAR protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	30kDa
Cellular Localization	Mitochondrion
Species Reactivity	Human
Positive Control	K-562 cells. Human adrenal or testicular carcinoma.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant StAR (Steroidogenic Acute Regulator) (Leydig Cell Marker) Antibody



Formalin-fixed, paraffin-embedded human testicular carcinoma stained with StARRecombinant Rabbit Monoclonal Antibody (STAR/3915R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified StARRecombinant Rabbit Monoclonal Antibody (STAR/3915R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Steroidogenic Acute Regulatory Protein (STAR) controls the rate-limiting step of steroidegenesis by translocating cholesterol from the outer mitochondrial membrane to the inner membrane where it is later cleaved to pregnenolone.It is primarily present in steroid-producing cells, including Leydig cells in the testis, theca cells and luteal cells in the ovary and adrenal cells in the adrenal cortex. Due to low levels of pregnenolone, seminomas and Leydig cell tumors display no specific STAR staining. Therefore, STAR antibody may assist in differentiating sex cord stromal tumors (SCST), seminomas and embryonal carcinomas.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.